

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

On the Infrared Properties of Dusty Torus

Liza Videla,¹ Paulina Lira,¹ Almudena Alonso-Herrero,² David M. Alexander,³ and Martin Ward³

Abstract. We performed imaging on 49 type 2 Seyfert galaxies in 6 near- and mid-infrared bands (1-10 μ m). We are separating the contribution of the torus from the host galaxy by radial profile fitting techniques and we will compare the observed spectral energy distributions with theoretical models of torus emission to constrain geometrical and physical parameters.

1. Introduction

In order to explain the differences observed between type 1 and type 2 AGN it has been proposed that an axially symmetric dusty structure (the torus) beyond the accretion disk absorbs a considerable fraction of the radiation emitted at wavelengths shorter than 1 μ m (the AGN Unified Scheme). The dust in this torus typically reaches temperatures of a few hundred degrees and therefore its emission peaks somewhere at IR wavelengths. It is therefore in this wavelength regime that the torus can be detected allowing us to determine its properties.

2. Observations, data analysis and preliminary results

We have obtained images for 49 Seyfert 2 galaxies in 6 IR bands: J (1.25 μ m), H (1.65 μ m), K (2.2 μ m), L (3.78 μ m), M (4.66 μ m) and N (10.36 μ m) using the VLT at Paranal, NTT at La Silla and Gemini at Pachón. These galaxies were selected from the *Extended 12 μ m Galaxy Sample* (Rush et al. 1993).

We obtained the surface brightness profiles of each galaxy and modeled them to separate the contribution from a nuclear point source and a stellar component (disk+bulge, Figure 1) to construct the nuclear spectral energy distributions (SEDs) for each galaxy. The SEDs are then fitted using theoretical models of the emission of the dusty torus from Nenkova, Ivezić, & Elitzur (2002), who solved the radiative transfer problem of a clumpy torus including absorption, emission and scattering.

In Figure 2 we show results for MCG -2-40-004 and MCG -3-34-64, both galaxies classified as type 2 Seyfert galaxies in Rush et al. (1993), but in Véron-Cetty & Véron (2006) MCG -2-40-004 is classified as a type 1.9 Seyfert, and MCG -3-34-64

¹Departamento de Astronomía, Universidad de Chile, Casilla 36D, Santiago, Chile

²Instituto Departamento de Astrofísica Molecular e Infrarroja, Instituto de Estructura de la Materia, CSIC, E-28006, Madrid, España

³Department of Physics, Durham University, South Road, Durham, DH1 3LE, UK.




Figure 1. Surface brightness profile of MCG -2-40-004 at H (left) and K (right) bands. Dots represent the data points, the dashed line the disk component, the dot-dashed line the bulge component, and the dotted line the nuclear contribution. The solid line is the sum of the 3 components.




Figure 2. Nuclear SEDs of MCG -2-40-004 (left) and MCG -3-34-64 (right), and their fitted model.

as a type 1 due to the detection of broad polarized Balmer lines. Following Nenkova et al. (2002) definitions, for MCG -2-40-004 we obtained a torus of 100 pc in size, with 5 clouds along the line of sight (each with a $\tau_\nu=20$) and an inclination of 60° . For MCG -3-34-64 we obtained a torus of 30 pc in size, with 5 clouds along the line of sight (each with a $\tau_\nu=40$) and a face on inclination. This SED is clearly more consistent with a type 1 Seyfert classification.

Complementary data for our study comes from X-ray, spectropolarimetry, ISOCAM and Spitzer observations. Together with our observations, we will use this information to further constrain the SEDs of our targets and determine the geometrical and physical parameters that govern the IR emission from AGNs.

Acknowledgments. LV thanks support by Conicyt to attend this meeting.

References

- Nenkova, M., Ivezić, Ž., & Elitzur, M. 2002, ApJ, 570, L9
 Rush, B., Malkan, M. A., & Spinoglio, L. 1993, ApJS, 89, 1
 Véron-Cetty, M.-P., & Véron, P. 2006, A&A, 455, 773